

KetoVie™ 4:1 (Low carbohydrate MCT enhanced lipid-based ketogenic formula)

PRODUCT INFORMATION

KetoVie 4:1 Chocolate 250 mL carton (8.5 fl. oz) Reimbursement Code: 24359-0501-03  
KetoVie 4:1 Vanilla 250 mL carton (8.5 fl. oz) Reimbursement Code: 24359-0502-03  
Manufactured by Cambrooke Therapeutics, Inc. Ayer, MA 01432 www.cambrooke.com

DISPENSE BY PRESCRIPTION

KetoVie 4:1 is a nutritionally complete medical food for the dietary management of intractable epilepsy, glucose transporter type 1 deficiency syndrome, pyruvate dehydrogenase deficiency and other disorders where a ketogenic diet is indicated.

DESCRIPTION

KetoVie 4:1 is a liquid ready to use prescription medical food for the clinical dietary management of intractable epilepsy, glucose transporter type 1 deficiency syndrome (Glut-1 DS), pyruvate dehydrogenase deficiency (PDH) and other disorders that require a ketogenic diet. KetoVie 4:1 is to be used only under medical supervision. KetoVie 4:1 has been developed, labeled and should be administered in accordance with the statutory and the Food and Drug Administration's (FDA's) regulatory definition of Medical Foods. Congress defined "Medical Food" in the Orphan Drug Act and Amendments of 1988 as a formulation to be administered enterally (for oral or tube feeding) under the supervision of a physician and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles are established by medical evaluation. 21 U.S.C. 360ee(b)(3). KetoVie 4:1 is supplied in a single dose, 8.5 fl oz (250 mL) cartons, 30 cartons per case. Each carton contains a 4:1 ketogenic ratio (4 grams of fat to every 1 gram of non-fat [protein and net carbohydrate\*]), 360 calories (Vanilla) and 390 calories (Chocolate), 100% whey protein source, 25% of total calories as medium chain triglyceride (MCT), and provides a complete micronutrient profile with enhanced amounts of selenium and carnitine. The caloric density (calories per mL) of KetoVie is 1.44 kcal/mL (Vanilla) and 1.56 kcal/mL (Chocolate).

\*The net carbohydrate weight (g) is the total carbohydrate minus the total dietary fiber.

PRIMARY INGREDIENTS

Ready to Use 4:1 Ketogenic Ratio

KetoVie 4:1 contains a 4:1 macronutrient ratio by weight of fat to combined protein and net carbohydrate\* weight. The classic 4:1 (ketogenic ratio) ketogenic diet is effective in the dietary management of intractable epilepsy<sup>1,2,3,4,5</sup> and is considered the diet of choice for the metabolic disorders glucose transporter type 1 deficiency syndrome (Glut-1 DS)<sup>6,7,8</sup> and pyruvate dehydrogenase deficiency (PDH).<sup>8,9</sup> Benefits of a liquid ready to use ketogenic medical food, such as KetoVie 4:1, whether as a sole source or supplemental nutrition, include palatability, ease of diet titration, reduced risk of calculation and preparation error, and improved compliance.<sup>10, 11</sup>

Fat and Medium Chain Triglycerides

KetoVie 4:1 contains a blend of saturated fats, polyunsaturated fats and monounsaturated fats. Refer to the Fat Profile Table below for a more detailed breakdown. KetoVie 4:1 has a saturated fat component with over 50% contribution from medium chain triglycerides (MCTs). MCTs are fats with 6-12 carbon length fatty acid chains. MCT can be beneficial to ketogenic diet management in that it is directly absorbed in the intestines and more efficiently processed into ketones in the liver compared to long chain triglycerides (LCTs). MCTs also have a laxative effect, which can be helpful for bowel regularity.<sup>12</sup> Algae-sourced long chain omega three fatty acid docosahexaenoic acid (DHA) is added at 76 milligrams per serving and KetoVie is additionally supplemented with essential fatty acids linoleic and α-linolenic acid.

FAT PROFILE			
	Measurement per 250mL serving	KetoVie 4:1 Vanilla	KetoVie 4:1 Chocolate
Total Fat	grams	35.4	38.4
Saturated Fat	grams	21.0	22.6
MCT	grams	12.0	12.0
Monounsaturated Fat	grams	10.9	11.9
Polyunsaturated Fat	grams	3.5	3.9
Linoleic Acid	milligrams	2250	2300
α-Linolenic Acid	milligrams	930	930
DHA	milligrams	76	76

Whey

KetoVie 4:1 protein source is from a 100% whey protein. Whey protein is a complete protein derived from milk. Cow and human milk contain both whey and casein protein fractions. Whey

protein has been shown to improve gastric emptying time in those with poor gastric motility compared to casein dominant milk proteins.<sup>13,14</sup> Impaired gastric motility can be common for those on a ketogenic diet.<sup>8</sup>

**Carnitine**

Carnitine is a conditionally essential amino acid necessary for transporting fatty acids into the cell's mitochondria for oxidation and is crucial for energy and ketone production. Carnitine deficiency is common among children and adolescents with epilepsy, especially among those receiving valproic acid therapy, a common medication used to treat seizures.<sup>15,16</sup> KetoVie 4:1 contains 50mg of carnitine per serving.

**Selenium**

Selenium is an essential trace element with many critical roles in the body. Selenium deficiency has been associated with ketogenic therapy. A serious complication of selenium deficiency is cardiomyopathy, which can be fatal.<sup>17</sup> Intakes of at least 20 mcg per day of selenium have been found to be protective against selenium deficiency related cardiomyopathy in adults.<sup>18</sup> KetoVie 4:1 contains 22 mcg of selenium per serving. The Reference Dietary Allowance (RDA) of selenium is 15-55 mcg per day depending on age with upper limits ranging between 45-400 mcg per day.<sup>18</sup>

**Fiber**

One of the most common complications related to ketogenic diet therapy is GI disturbance, involving nausea/vomiting, diarrhea, or constipation. Because ketogenic diets restrict carbohydrates, it may be difficult to meet dietary fiber recommendations. KetoVie 4:1 contains a blend of soluble and insoluble fiber which can be helpful for bowel regularity and may help to maintain healthy cholesterol levels.<sup>19</sup> Sources of KetoVie 4:1 fiber include inulin, cellulose, and pectin.

Dietary Fibers	Chocolate (grams)	Vanilla (grams)
Total	5.2	4.4
Soluble	3.9	3.3
Insoluble	1.3	1.1

**Micronutrients and Macronutrients**

Patients utilizing ketogenic therapy follow a severely restricted diet to minimize carbohydrate intake and provide the necessary fats at the desired ketogenic ratio for the production of ketone bodies. As such, there is meaningful risk and challenges in receiving recommended daily intake of many nutrients.<sup>20</sup> Many commercially available vitamin and mineral supplements also contain carbohydrate which must be factored into the diet. To compensate for this, KetoVie 4:1 includes a full profile of micronutrients and macronutrients, as well as the enhanced levels of selenium and carnitine as noted above. With a documented potential risk of bone fractures<sup>21</sup> when on ketogenic therapy along with the challenge of meeting recommended micronutrients noted above, KetoVie's formulation contains a special blend of vitamins and minerals important for bone health including: Vitamin D3, calcium, phosphorus, magnesium, and Vitamin K2 - menaquinone-7.

The use of oral citrates has been shown to be an effective preventative supplement against kidney stones (renal calculi) in children who receive the ketogenic diet.<sup>22</sup> KetoVie 4:1 contains citrates (potassium citrate and sodium citrate) at 26 mEq (Vanilla) and 30 mEq (Chocolate) per serving.

**Sugar Substitute**

KetoVie 4:1 Vanilla is sweetened with a blend of sucralose and acesulfame potassium, sugar substitutes that contribute little to the carbohydrate total and do not add significant calories. KetoVie 4:1 Chocolate is sweetened with sucralose only. Both KetoVie 4:1 Vanilla and Chocolate flavors are aspartame free.

**Complete Ingredients**

Water, cocoa butter, medium chain triglycerides, canola oil, whey protein blend (whey protein concentrate and whey protein isolate) (milk), inulin, fortification blend (tricalcium phosphate, choline bitartrate, sodium ascorbate and ascorbic acid, magnesium phosphate, L carnitine l-tartrate, niacinamide, dl-alpha-tocopheryl acetate, zinc sulfate, calcium d-pantothenate, manganese sulfate, iron, riboflavin, pyridoxine HCl, thiamin HCl, folic acid, vitamin A palmitate, potassium iodide, copper, sodium selenite, sodium molybdate, K1 phytonadione, K2 menaquinone-7, chromium chloride, biotin, vitamin D3, cyanocobalamin, maltodextrin), natural flavors, cellulose gel, pectin, potassium citrate, sodium hexametaphosphate, sodium citrate, sodium steryl lactylate (soy), DHA algal oil, salt, carrageenan, L-histidine, sucralose. Contains milk and soy. Also contains corn.

KetoVie 4:1 Vanilla also contains acesulfame potassium.  
KetoVie 4:1 Chocolate also contains cocoa processed with alkali.

*Low in lactose (0.2g/250mL serving) and aspartame free.*

**GENERALLY RECOGNIZED AS SAFE**

The ingredients in KetoVie 4:1 are Generally Recognized As Safe (GRAS). This is the statutory safety standard of the U.S. Food and Drug Administration (FDA). The use of an ingredient may be established as GRAS based on documented widespread common use of the ingredient in foods prior to January 1, 1958, or based on scientific procedures. GRAS status based on scientific

procedures requires technical evidence of safety (i.e., a reasonable certainty of no harm under the conditions of intended use), and evidence of general recognition of such safety amongst qualified experts.

**MEDICAL FOOD STATUS**

KetoVie 4:1 has been developed, labeled and should be administered in accordance with FDA statutory and regulatory definition of Medical Foods. Congress defines "Medical Food" in the Orphan Drug Act and Amendments of 1988 as a formulation to be administered enterally (for oral or tube feeding) under the supervision of a physician and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles are established by medical evaluation.

**INDICATIONS FOR USE**

For the dietary management of intractable epilepsy, glucose transporter type 1 deficiency syndrome (Glut-1 DS), pyruvate dehydrogenase deficiency (PDH) and other disorders where a ketogenic diet is indicated. KetoVie 4:1 should always be used under medical supervision. KetoVie 4:1 may be used as sole source nutrition and as a supplement.

AMOUNT OF KETOVIE 4:1 NEEDED TO MEET 100% MICRONUTRIENT NEEDS*	
Age (years)	Amount
1-3	700 mL (1,000 kcal)
4-8	1,000 mL (1,440 kcal)
Older children and adults	Varies based on age and gender

\*Excludes electrolytes

**CLINICAL EXPERIENCE**

Ketogenic diet therapy has been used in the treatment of seizures since 1921. There has been a resurgence in its use in recent years.<sup>23</sup> Several recent meta-analysis reviews have looked at ketogenic diet therapy and have shown that ketogenic diet therapy is an effective treatment in reducing the number of seizures in children with intractable seizures, though not every patient will respond.<sup>4,5,24</sup> One of the meta analysis studies pooled data from over 1,084 pediatric epileptic patients and found the ratio of treatment success (>50% seizure reduction) to diet discontinuation was 2.25 (95% confidence interval 1.69-2.98), indicating a high level of efficacy for ketogenic diet therapy and seizure reduction.<sup>5</sup> A 2008 randomized control trial (Neal, et al.) of 145 children and adolescents with intractable epilepsy found 38% of patients achieved >50% reduction of seizures after 3 months of ketogenic diet therapy compared to controls. Another 7% achieved >90% reduction in seizures.<sup>1</sup> Adolescent and adult data indicate ketogenic diet therapy to be equally efficacious in these groups.<sup>2,4</sup> There are studies showing benefit of ketogenic diet therapy to several different seizure types and other conditions.<sup>8,25</sup>

**Conditions Shown to Respond to Ketogenic Diet Therapy**

- Glucose transporter protein 1 (GLUT-1) deficiency
  - Pyruvate dehydrogenase deficiency (PDHD)
  - Myoclonic-astatic epilepsy (Doose syndrome)
  - Tuberous sclerosis complex
  - Rett syndrome
  - Severe myoclonic epilepsy of infancy (Dravet syndrome)
  - Lennox Gastaut syndrome
- Infantile spasms
  - Selected mitochondrial disorders
  - Glycogenosis type V
  - Landau-Kleffner syndrome
  - Lafora body disease
  - Subacute sclerosing panencephalitis (SSPE)

**PHARMACOKINETICS**

Ketogenic diet therapy is meant to put the body into a state of ketosis by using ketone bodies for energy. Ketone bodies are the by-product of fat metabolism and can be used as an alternative to glucose as a fuel source for the brain. Ketosis is brought about by the oxidation of body fat stores, such as during times of prolonged fasting, or when the body is fed a diet primarily made up of fats. KetoVie 4:1 is a specially formulated ketogenic medical food at a classic 4:1 ketogenic ratio (4 grams of fat to every 1 gram of non-fat [protein and net carbohydrate.]) This 4:1 ketogenic ratio has been found to be effective in achieving the desired level of ketosis.<sup>8</sup> Fats are metabolized in a metabolic pathway known as beta oxidation. In this multi-step process, fatty acids are taken up by cells and transported into the mitochondria by a carnitine shuttle, hence carnitine's important role in fatty acid metabolism. In the mitochondria, the activated fatty acids are converted into acetyl-CoA. Acetyl-CoA can then be further processed into energy via the tricarboxylic acid cycle (TCA cycle) or taken to the liver to be turned into ketone bodies. Ketone bodies can pass through the blood brain barrier and may play a role in ketogenic diet therapy's suppression of seizure activity in the brain. Included in the KetoVie 4:1 formulations are 25% of calories from medium chain triglycerides (MCTs.) MCTs are fats with 6-12 carbon length fatty acid chains. MCTs are particularly beneficial to achieving the desired level of ketosis in the body as they are absorbed directly from the intestines and sent directly to the liver for efficient processing into ketone bodies.<sup>12</sup>

**Precautions and Contraindications**

KetoVie 4:1 contains protein from whey, a milk protein. Therefore, it may not be suitable for those with an allergy to milk or milk products. There are certain conditions in which a ketogenic diet may be contraindicated. These may include but are not limited to the conditions listed below.<sup>8</sup>

## Conditions Contraindicated to Use of Ketogenic Diet Therapy

- Carnitine deficiency (primary)
- Carnitine palmitoyltransferase (CPT) I or II deficiency
- Carnitine translocase deficiency
- Beta-oxidation defects
- Medium-chain acyl dehydrogenase deficiency (MCADD)
- Long-chain acyl dehydrogenase deficiency (LCADD)
- Short-chain acyl dehydrogenase deficiency (SCADD)
- Long-chain 3-hydroxyacyl-CoA deficiency
- Medium-chain 3-hydroxyacyl-CoA deficiency
- Pyruvate carboxylase deficiency
- Porphyria
- Inability to maintain adequate nutrition
- Noncompliance

## Adverse Reactions

There are no known incremental adverse reactions for the use of KetoVie 4:1 as a part of a ketogenic diet. Ketogenic diets have been associated with certain adverse reactions. These can include metabolic abnormalities, gastrointestinal distress (vomiting, constipation, diarrhea, abdominal discomfort,) secondary carnitine deficiency, hypercholesterolemia, renal calculi, delayed linear growth in children and increased risk of bone fractures.<sup>3,25,26,27,28, 29</sup> Close medical supervision is required.

## Drug Interactions

KetoVie 4:1 when prescribed as supplemental or sole source nutrition may impact how some drugs are metabolized. All medications should be discussed with a physician or pharmacist. Medical supervision by a physician is required.

## Toxicity

None known.

## SPECIAL POPULATIONS

KetoVie 4:1 is indicated for patients 12 months and older requiring a ketogenic diet. Always check with physician for proper dosage recommendations.

## DOSAGE AND ADMINISTRATION

Use as directed by physician. Must be administered under medical supervision only. Ready to drink. Shake well. Store in a cool, dry place. Refrigerate after opening and store refrigerated up to 24 hours. Once opened, product should be kept for no longer than 4 hours at room temperature. Do not freeze.

## HOW SUPPLIED

KetoVie 4:1 is supplied in a single dose, 8.5 fl oz (250 mL) cartons, 30 cartons per case. Keep in a cool, dry place until ready to use. Reimbursement Code: Vanilla: 24359-0502-03, Chocolate: 24359-0501-03.

## REFERENCES

1. Neal EG, Chaffé H, Schwartz RH, Lawson MS, Edwards N, Fitzsimmons G, Whitney A, Cross JH. The ketogenic diet for the treatment of childhood epilepsy: a randomised controlled trial. *Lancet Neurol*. 2008 Jun;7(6):500-6.
2. Payne NE, Cross JH, Sander JW, Sisodiya SM. The ketogenic and related diets in adolescents and adults—a review. *Epilepsia*. 2011 Nov;52(11):1941-8.
3. Whless JW. The ketogenic diet: an effective medical therapy with side effects. *J Child Neurol*. 2001 Sep;16(9):633-5.
4. Ye F, Li XJ, Jiang WL, Sun HB, Liu J. Efficacy of and patient compliance with a ketogenic diet in adults with intractable epilepsy: a meta-analysis. *J Clin Neurol*. 2015 Jan;11(1):26-31.
5. Henderson CB, Filloux FM, Alder SC, Lyon JL, Caplin DA. Efficacy of the ketogenic diet as a treatment option for epilepsy: meta-analysis. *J Child Neurol*. 2006 Mar;21(3):193-8.
6. Klepper J. Impaired glucose transport into the brain: the expanding spectrum of glucose transporter type 1 deficiency syndrome. *Curr Opin Neurol*. 2004 Apr;17(2):193-6.
7. Klepper J, Voit T. Facilitated glucose transporter protein type 1 (GLUT1) deficiency syndrome: impaired glucose transport into brain—a review. *Eur J Pediatr*. 2002 Jun;161(6):295-304.
8. Kossoff EH, Zupec-Kania BA, Amark PE, Ballaban-Gil KR, Christina Bergqvist AG, Blackford R, Buchhalter JR, Caballo RH, Helen Cross J, Dahlin MG, Donner EJ, Klepper J, Jehle RS, Kim HD, Christiansa Liu YM, Nattion J, Nordli DR Jr, Pfeiffer HH, Rho JM, Stafstrom CE, Thiele EA, Turner Z, Wirrell EC, Whless JW, Veggiotti P, Vining EP, Charlie Foundation, Practice Committee of the Child Neurology Society; Practice Committee of the Child Neurology Society; International Ketogenic Diet Study Group. Optimal clinical management of children receiving the ketogenic diet: recommendations of the International Ketogenic Diet Study Group. *Epilepsia*. 2009 Feb;50(2):304-17.
9. Wexler ID, Hemalatha SG, McConnell J, Buist NR, Dahl HH, Berry SA, Cederbaum SD, Patel MS, Kerr DS. Outcome of pyruvate dehydrogenase deficiency treated with ketogenic diets. *Studies in patients with identical mutations. Neurology*. 1997 Dec;49(6):1655-61.
10. Kossoff EH, McGrogan JR, Freeman JM. Benefits of an all-liquid ketogenic diet. *Epilepsia*. 2004 Sep;45(9):1163.
11. Hosain SA, La Vega-Talbot M, Solomon GE. Ketogenic diet in pediatric epilepsy patients with gastrostomy feeding. *Pediatr Neurol*. 2005 Feb;32(2):81-3.
12. Marten B, Pfeuffer M, Schrezenmeir J. Medium-chain triglycerides. *Int Dairy J*. 2006 Jun;16: 1374-1382.
13. Savage K, Kritis S, Schwarzer A, Davidson G, Omar T. Whey- vs casein-based enteral formula and gastrointestinal function in children with cerebral palsy. *JPEN J Parenter Enteral Nutr*. 2012 Jan;36(1 Suppl):118S-23S.
14. Kuyumcu S, Menne D, Curcic J, Goetze O, Klebach M, Abrahamse E, Hofman Z, Fried M, Schwizer W, Steingotter A. A Noncoagulating Enteral Formula Can Empty Faster From the Stomach: A Double-Blind, Randomized Crossover Trial Using Magnetic Resonance Imaging. *JPEN J Parenter Enteral Nutr*. 2014 Apr 3. [Epub ahead of print]
15. Raskind JY, El-Chaar GM. The role of carnitine supplementation during valproic acid therapy. *Ann Pharmacother*. 2000 May;34(5):630-8.
16. De Vivo DC, Bohan TP, Coulter DL, Dreifuss FE, Greenwood RS, Nordli DR Jr, Shields WD, Stafstrom CE, Tein I. L-carnitine supplementation in childhood epilepsy: current perspectives. *Epilepsia*. 1998 Nov;39(11):1216-25.
17. Sirikonda NS, Patten WD, Phillips JR, Mullett CJ. Ketogenic diet: rapid onset of selenium deficiency-induced cardiac decompensation. *Pediatr Cardiol*. 2012 Jun;33(5):834-8.
18. National Institutes of Health. Selenium: Dietary Supplement Fact Sheet. [Webpage]. Available at: <http://ods.od.nih.gov/factsheets/Selenium-HealthProfessional/>. Accessed: Feb 03, 2015.
19. Anderson JW, Baird P, Davis RH Jr, Ferreri S, Knudtson M, Koraym A, Waters V, Williams CL. Health benefits of dietary fiber. *Nutr Rev*. 2009 Apr;67(4):188-205.
20. Zupec-Kania B, Zupanc ML. Long-term management of the ketogenic diet: seizure monitoring, nutrition, and supplementation. *Epilepsia*. 2008 Nov;49 Suppl 8:23-6.
21. Kang HC, Chung DE, Kim DW, Kim HD. Early- and Late-onset Complications of the Ketogenic Diet for Intractable Epilepsy. *Epilepsia*. 2004; 45(9):1116-1123.
22. McNally MA, Pzyk PL, Rubenstein JE, Hamdy RF, Kossoff EH. Empiric use of potassium citrate reduces kidney-stone incidence with the ketogenic diet. *Pediatrics*. 2009 Aug;124(2):e300-4.
23. Kossoff EH. More fat and fewer seizures: dietary therapies for epilepsy. *Lancet Neurol*. 2004 Jul;3(7):415-20.
24. Li HF, Zou Y, Ding G. Therapeutic Success of the Ketogenic Diet as a Treatment Option for Epilepsy: a Meta-analysis. *Iran J Pediatr*. 2013 Dec;23(6):613-20.
25. Caraballo RH, Fortini S, Fresler A, Armeno M, Anela A, Cresta A, Mestre G, Escobal N. Ketogenic diet in patients with Lennox-Gastaut syndrome. *Seizure*. 2014 Oct;23(9):751-5.
26. Vining EP, Pzyk P, McGrogan J, Hladky H, Anand A, Krieger S, Freeman JM. Growth of children on the ketogenic diet. *Dev Med Child Neurol*. 2002 Dec;44(12):796-802.
27. Peterson SJ, Tangney CC, Pimentel-Zablah EM, Hjelmgren B, Booth G, Berry-Kravis E. Changes in growth and seizure reduction in children on the ketogenic diet as a treatment for intractable epilepsy. *J Am Diet Assoc*. 2005 May;105(5):718-25.
28. Ballaban-Gil K, Callahan C, O'Dell C, Pappo M, Moshé S, Shinnar S. Complications of the ketogenic diet. *Epilepsia*. 1998 Jul;39(7):744-8.
29. Liu YM, Williams S, Basualdo-Hammond C, Stephens D, Curtis R. A prospective study: growth and nutritional status of children treated with the ketogenic diet. *J Am Diet Assoc*. 2003 Jun;103(6):707-12.